The Structure and Dynamics of Macrozoobenthic Communities of the Bega River Water

Anca-Andreea MARIN^{1*}, Benoni LIXANDRU¹, Milca PETROVICI², Adrian SINITEAN², Florica MORARIU¹, Dumitru POPESCU¹

¹Banat University of Agricultural Sciences and Veterinary Medicine “King Michael I of Romania” from Timisoara, Faculty of Animal Science and Biotechnology,

²West University of Timișoara, Faculty of Chemistry Biology and Geography, Department of Chemistry-Biology, Timișoara, Romania

*corresponding author, e-mail: marin.aandreea@yahoo.com

Bulletin UASVM series Agriculture 73(1)/2016
Print ISSN 1843-5246; Electronic ISSN 1843-5386
DOI 10.15835/buasvmcn-agr: 12010

ABSTRACT

The benthic macro invertebrates are good indicators of localized conditions, as many of the benthic macro invertebrates have limited migration patterns or a sessile mode of life. Most species have a complex life cycle of one year or more. Sensitive life stages will respond quickly to stress; the overall community will respond more slowly. The structure and distribution of the different benthonic macro-invertebrates communities is considered to be one of the important indicators of the surface waters' quality. In November 2014, were collected 20 quantitative samples of benthic zones, in the Bega River in order to highlight the links between pollution degree and the saprobionte organisms community. The aim of this paper is to show the effects of human impact (Ghiroda village, potable water treatment station, sewage water treatment station of Timisoara, Sânmihaiu Roman village) on benthic community structure and dynamics of different ecosystems from Bega River water. After the laboratory work was carried out, it was identified ten groups of benthic macro invertebrates: *Oligochaeta* subclass *Hirudin* class, *Lamelibranchiata* class, *Gastropoda* class, *Nematoda* phylum, Diptera order (larvae of the families *Chironomidae*, *Ceratopogonidae* and *Tipulidae*), *Isopoda* order, *Trichoptera* order, *Odonata* order, *Coleoptera* order).

Keywords: *Bega river, bentonic water quality, macro invertebrates.*

INTRODUCTION

An important indicator of water quality in aquatic ecosystems is the structure of macrozoobenthos communities (Rosenberg & Resh, 1993). Benthic invertebrates are sensitive indicators of ecosystem quality change, because they highlight the long-term changes in water quality, unlike assessments physical-chemical factors that highlight situations of moment (Johnson *et al.*, 1993).

The structure and distribution of the different benthonic macro-invertebrates communities is considered to be one of the important indicators of the surface waters' quality (Badea *et al.*, 2010; Kubosova *et al.*, 2010; Ogbeibu & Oribhabor, 2002).

Benthic macro invertebrates have a fundamental role in absorbing and filtering substances in aquatic ecosystems, reason for that they are good indicators of organic load, in the main, and of mineral water pollution.

Benthic invertebrates fauna is considered one constant compartment in the structure of the biocenosis water systems (Botnariuc & Vădineanu 1982; Botnariuc *et al.*, 1987; Ignat, 1986; Chaloner *et al.*, 2009; Cooper *et al.*, 2009 Wallace & Eggert, 2009; Whiles & Grubaugh, 2000).

The benthic macro invertebrates are good indicators of localized conditions, as many of the benthic macro invertebrates have limited migration patterns or a sessile mode of life. Most

species have a complex life cycle of one year or more. Sensitive life stages will respond quickly to stress; the overall community will respond more slowly (Moldoveanu & Rîșnoveanu, 2010).

The aim of this paper is to show the effects of human impact (Ghiroda village, potable water treatment station, sewage water treatment station of Timisoara, Sânmihaiu Roman village) on benthic community structure and dynamics of Bega River water.

MATERIAL AND METHODS

The samples have been collected in November 2014, when were collected 20 quantitative samples of benthic zones, in the Bega River in order to show the effects of human impact on benthic community structure and dynamics of different ecosystems from Bega River water.

Samples were collected from the upstream, middle and downstream of Timisoara city. The benthic samples were collected with Ekman-sampler with a surface of 225 cm² and were subsequently washed with benthic nets (meshes of 250 µm) and stored in 8% formaldehyde (Marin *et al.*, 2014; Lixandru, 2006; Péterfi & Sinitean, 2002; Petrovici, 2009).

The collecting stations (S) were located as follows (figure 1):

S1 is located upstream of Timisoara city, near Ghiroda village, upstream of potable water treatment station.

S2 is located upstream of sewage water treatment station of Timisoara.

S3 is located near Sânmihaiu Roman village from Timis County and downstream of sewage water treatment station of Timisoara.

S4 is located near Otelec village, before the border line with Serbia Country.

There have been calculated the density ($D_i = n_i / S_p$), the abundance ($A = (n_i / N) * 100$) and the frequency ($F = N_i * 100 / N_p$), where n_i represents the total number of individuals for the i series, S_p the total researched area, N the total number of individuals belonging to all species (from the sample or the studied samples), N_i the number of stations within which been identified the subjected species, N_p the total number of stations (Sîrbu & Benedek, 2004; Stan, 1995).

RESULTS AND DISCUSSION

After the laboratory work was carried out, it was identified ten groups of benthic macroinvertebrates (table 1): *Oligochaeta* subclass *Hirudin* class, *Lamelibranchiata* class, *Gastropoda* class, *Nematoda* phylum, Diptera order (larvae of the families *Chironomidae*, *Ceratopogonidae* and *Tipulidae*), *Isopoda* order, *Trichoptera* order, *Odonata* order, *Coleoptera* order and macro invertebrates density is presented in figure 2.

The high tolerance to the impurification of *Oligochaeta* subclass and *Chironomidae* order has been demonstrated in numerous studies

Fig. 1. The location of the sample collecting stations on Bega River water

(Benbow, 2009; Collier *et al.*, 2010; Courtney & Merritt, 2009; Marchese *et al.*, 2008). These two groups of invertebrates show significantly larger tolerance limits, adapting to various environmental conditions (Lucan-Bouché *et al.*, 1999; Verdonschet, 1999).

At the first station (S1) have been identified groups which Lorenz (2003) considered that they are indicators of unpolluted water (*Lamelibranchiata* class, *Gastropoda* class, *Odonata* order).

Analyzing the macro invertebrates density (fig. 2), we can say that at the first stations the density

of individual's that belong to the *Oligochaeta* subclass and *Chironomidae* order have a density which is smaller than the density of individual's that belong to the *Gastropoda* class which means that the degree of impurification of this station is very small.

Analyzing the macro invertebrates density at the second station (fig. 2) we can notice an increase of the density values of groups which are classified as indicators of unpolluted water (*Lamelibranchiata* class, *Gastropoda* class). At this station we can notice the highest density values of sensitive groups to the pollution

Fig. 2. Macro invertebrate's density (individual's m²) from Bega River

Tab. 1. Groups of saprobionți in relation with the collection stations

Groups	Station 1 (S1)	Station 2 (S2)	Station 3 (S3)	Station 4 (S4)
<i>Oligochaeta</i>	x	x	x	x
<i>Hirudinea</i>			x	
<i>Lamelibranchiata</i>	x	x		
<i>Gastropoda</i>	x	x		x
<i>Nematoda</i>	x	x		x
<i>Chironomidae</i>	x	x	x	x
<i>Ceratopogonidae</i>	x	x		x
<i>Tipulidae</i>			x	x
<i>Isopoda</i>			x	
Trichoptera				x
<i>Odonata</i>	x			
<i>Coleoptera</i>	x			

X= the presence

At the station three (S3) we can observe that the values of density for sensitive groups to the pollution (*Lamelibranchiata* class, *Gastropoda* class) start to decline and increase the density values of groups show significantly larger tolerance limits of pollution (*Oligochaeta* subclass and *Chironomidae* order).

At this station we can notice the highest density values to the groups who have a high tolerance to the pollution.

Also at S3 are identified individual's that belong to the *Hirudin* class and *Isopoda* order, groups who show significantly larger tolerance limits at impurification and the disappear the individual's that belong to the *Lamelibranchiata* class, *Gastropoda* class, *Odonata* order.

The last station (S4) it is located near Otelec village, before the border line with Serbia Country. At this station decrease the density values of individual's that belong to the *Oligochaeta* subclass and *Chironomidae* order and appear the individual's that belong to the *Gastropoda* class. Also at this station are identified individual's than belong to the Trichoptera order, macro invertebrates who are considered indicators of the indicators of unpolluted water (Lorenz, 2003).

Regarding to the numerical abundance we can notice that is in correlation with the density, if the values of density grow up, then the numerical abundance shows increased values (fig. 3).

Analysing the frequency (fig. 4), individual's belong to the *Oligochaeta* subclass show a 80 % frequency at the first station, a 60 % frequency at the second station and a 100% frequency at the three and the last station.

Individual's belong to the *Diptera* order (larvae of the *Chironomidae* families) show a 60 % frequency at the first station, a 80 % frequency at the second station and a 100% frequency at the three and the last station.

Individual's belong to the *Gastropoda* class show a 100 % frequency at the first station, a 80 % frequency at the second station and a 20 % frequency at the last station (fig. 4).

Macro invertebrates belong to the *Lamelibranchiata* class has show a 100 % frequency at the first station, a 80 % frequency at the second station and at station 3 and 4 the this invertebrates disappear.

Individual's belong to the *Trichoptera* order has a 40 % frequency at the last station.

Macro invertebrates belong to the *Nematoda* phylum show a 20 % frequency at the first and the second station, individual's belong to the *Hirudin* class has a 60 % frequency at the station three and individual's belong to the *Isopoda* order has a 20 % frequency at the same station (fig. 4).

Individual's belong to the *Odonata* order show a 20 % frequency at station 1.

Fig. 3. The numerical abundance of the invertebrates group at: a- first station, b- station 2, c- station 3, d- station 4.

Fig. 4. Macro invertebrate's frequency in the Bega River (%)

CONCLUSIONS

Have been identified ten groups of benthic macro invertebrates: *Oligochaeta* subclass *Hirudin* class, *Lamelibranchiata* class, *Gastropoda* class, *Nematoda* phylum, *Diptera* order (larvae of the families *Chironomidae*, *Ceratopogonidae* and *Tipulidae*), *Isopoda* order, *Trichoptera* order, *Odonata* order, *Coleoptera* order.

In conclusions we can say:

at the first and the second station has been identified groups who are considered indicators of unpolluted water (*Lamelibranchiata* class, *Gastropoda* class, *Odonata* order), and at the three station this groups disappear,

at the second station was notice the highest density values to the groups who are indicators of unpolluted water and to the three station was notice the highest density values to the groups who have a high tolerance to the pollution (*Oligochaeta* subclass, *Diptera* order),

at the last station decrease the density values of the groups who have a high tolerance to the pollution and appear macro invertebrates who are considered indicators of the indicators of unpolluted water.

Acknowledgments. On this way I want to thank my scientific mentor and all my collaborators for their support in achieving these experiments.

REFERENCES

1. Badea, B. A., Gagy-Palfy, A., Stoian, L. C., Stan, G., (2010). Preliminary studies of quality assessment of aquatic environments from Cluj suburban areas, based on some

invertebrates bioindicators and chemical indicators. *AACL Bioflux* 3(1):35-41.

2. Benbow, M. E., (2009). Annelida, Oligochaeta and Polychaeta. *Encyclopedia of Inland Waters* 55:124-127.
3. Botnariuc, N., Vădineanu, A. (1982). *Ecologie*. Editura Didactică și Pedagogică, București.
4. Botnariuc, N., Ignat, Gh., Diaconu, I., Vădineanu, A. (1987). L'évolution de l'état trophique des écosystèmes aquatiques caractéristiques au Delta du Danube: 5. La structure et la dynamique de la faune benthonique, *Rev. Roum. Biol. Anim.* 32 (2): 121-127.
5. Chaloner, D.T., Hershey, A.E., Lamberti, G.A. (2009), *Benthic Invertebrate Fauna*. In: Editor-in-Chief: Gene E. Likens, Editor(s)-in-Chief, *Encyclopedia of Inland Waters*, Academic Press, Oxford, pp. 157-172.
6. Collier, K. J., Winterbourn, M. J., Jackson, R. J. (2010). Impacts of wetland afforestation on the distribution of benthic invertebrates in acid streams of Westland, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 23:479-490.
7. Cooper Z.N., McGuire A., Jones S., Le Grand J. (2009) *Equity, waiting times, and NHS reforms: retrospective study*. *BMJ* 339: b3264 doi10.1136/bmj/b3264.
8. Courtney, G. W., Merritt, R. W. (2009). *Diptera (Non-Biting Flies)*. *Encyclopedia of Inland Waters* 55:288-298.
9. Ignat, G. (1986). *Ecologia unor populatii de chironomide din apele Deltei Dunarii*. PhD thesis. University of Bucharest.
10. Johnson, M. K., Hashtroudi, S., & Lindsay, D. S. (1993). Source monitoring. *Psychological Bulletin*, 114, 3-28.
11. Kubosova, K., Brabec, K., Jarkovsky, J., Syrovatka, V. (2010). Selection of indicative taxa for river habitats: a case study on benthic macroinvertebrates using indicator species analysis and the random forest methods. *Hydrobiologia* 651:101-114.
12. Lixandru, B. (2006). *Metodologie ecologică*. Editura Eurobit, Timișoara.

13. Lorenz, C. M. (2003). Bioindicators for ecosystem management, with special reference to freshwater systems. *Trace Metals and other Contaminants in the Environment* 6:123-152.
14. Lucan-Bouché, M. L., Risbourg, S. B., Arsac, F., Vernet, G. (1999). An original decontamination process developed by the aquatic oligochaete *Tubifex tubifex* exposed to copper and lead. *Aquatic Toxicology* 45:19-17.
15. Marchese, M. R., Rodriguez, A. R., Pave, P. J., Carignano, M. R. (2008). Benthic invertebrates structure in wetlands of a tributary of the middle Parana River (Argentina) affected by hydrologic and anthropogenic disturbances. *Journal of Environmental Biology* 29(3): 343-348
16. Marin, Anca-Andreea, Lixandru, B., Petrovici, Milca, Sinitean A., Morariu Florica, Popescu D. (2014). Research regarding the diversity of the saprobionte categories in Bega river waters, Book of abstracts, No 1/ 2014, USAMV Cluj –Napoca,
17. Moldoveanu, M., Rîsnoveanu, G. (2010). Achievements and limitations for monitoring and assessing ecological integrity of lotic systems, National Conference NIHWM, 28-30 September, Bucharest, Cd version: 236-244.
18. Ogbeibu, A. E., Oribhabor, B. J. (2002). Ecological impact of river impoundment using benthic macro-invertebrates as indicators. *Water Research* 36:2427-2436.
19. Sîrbu, I., Benedek, A., M., (2004). *Practical Ecology*, Editura Universității Lucian Blaga, pp. 28-254, Sibiu,
20. Péterfi, L.Șt. & Sinitean, A. (2002). Preliminary Studies on Diatoms in Thermo mineral Springs from Băile Herculane (Caraș – Severin District), *Studies of Biodiversity, West Romania Protected Areas, USAMVBT*, 25-28.
21. Petrovici, Milca, (2009). Evaluarea calității apei râului Crișul Repede utilizând larvele de efemeroptere (Insecta: Ephemeroptera) ca bioindicatori), Editura Universității din Oradea
22. Rosenberg, D. M. and V. H. Resh. (1993). Introduction to freshwater biomonitoring and benthic macroinvertebrates. In: D.M. Rosenberg and V.H. Resh, eds. *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York.
23. Stan, G., (1995). Statistical methods with applications in entomological research (VII)]. *Bul Inf Soc Lepid Rom* 6:67-96. [In Romanian]
24. Verdonchet, P. F. M. (1999). Microdistribution of oligochaetes in a soft-bottomed lowland stream. *Hydrobiologia* 139:149-163.
25. Wallace, J.B. & Eggert, S.L. (2009). Benthic Invertebrate Fauna, Small Streams. In: Editor-in-Chief: Gene E. Likens, Editor(s)-in-Chief, *Encyclopedia of Inland Waters*, Academic Press, Oxford, pp. 173-190.
26. Whiles, M.R., Grubaugh, J.W., (2009). Benthic Invertebrate Fauna, River and Floodplain Ecosystems. In: Editor-in-Chief: Gene E. Likens, Editor(s)-in-Chief, *Encyclopedia of Inland Waters*, Academic Press, Oxford, pp. 205-215